


North Carolina State Health Director's Standing Order for Naloxone

This standing order signed by the North Carolina State Health Director authorizes any pharmacist practicing in the state of North Carolina and licensed by the North Carolina Board of Pharmacy to dispense the following Naloxone products to persons as directed below.

Naloxone HCl Dispensing Protocol			
Eligible Candidates	<ul style="list-style-type: none"> ▪ Persons who voluntarily request Naloxone and are at risk of experiencing an opiate-related overdose, including, but not limited to: <ul style="list-style-type: none"> – Current illicit or non-medical opioid users or persons with a history of such use – Persons with a history of opioid intoxication or overdose and/or recipients of emergency medical care for acute opioid poisoning – Persons with a high dose opioid prescription (>50 morphine mg equivalents per day) – Persons with an opioid prescription and known or suspected concurrent alcohol use – Persons from opioid detoxification and mandatory abstinence programs – Persons entering methadone maintenance treatment programs (for addiction or pain) – Persons with opioid prescription and smoking/COPD or other respiratory illness or obstruction – Persons with an opioid prescription who also suffer from renal dysfunction, hepatic disease, cardiac disease, HIV/AIDS – Persons who may have difficulty accessing emergency medical services – Persons enrolled in prescription lock in programs ▪ Persons who voluntarily request Naloxone and are the family member or friend of a person at risk of experiencing an opiate-related overdose. ▪ Persons who voluntarily request Naloxone and are in the position to assist a person at risk of experiencing an opiate-related overdose. 		
Route(s) of Administration	Intranasal (IN) <i>Preferred method</i>		Intramuscular (IM) Inject into shoulder or thigh
Medication and Required Device for Administration	Naloxone HCl 1 mg/mL Inj. 2 x 2 mL as pre-filled Luer-Lock syringes <ul style="list-style-type: none"> ▪ Dispense 2 (two) doses 2 (two) x Intranasal Mucosal Atomizing Devices (MAD 300) Available from: Teleflex (866-246-6990) or Safety Works, Inc. (800-723-3892)	Narcan @ 4 mg/0.1 mL Nasal Spray <ul style="list-style-type: none"> ▪ Dispense 1 x two-pack 	<u>Naloxone HCl 0.4mg/mL Inj.</u> <ul style="list-style-type: none"> ▪ 2 x 1mL single dose vials (SDV) ▪ 2 (two) 3 mL syringe ▪ 2 (two) 25 G, 1 inch needle <u>Naloxone HCl 2 mg/2mL Inj.</u> <ul style="list-style-type: none"> ▪ Dispense 2 (two) pre-filled syringes ▪ 2 (two) 25 G, 1 inch needle
Directions for Use	Call 911. Spray 1 mL in each nostril. Repeat every 3 minutes as needed if no or minimal response.	Call 911. Administer a single spray of NARCAN® in one nostril. Repeat every 3 minutes as needed if no or minimal response.	Call 911. Inject the entire solution of the vial or pre-filled syringe IM in shoulder or thigh. Repeat every 3 minutes as needed if no or minimal response.
Refills	PRN		
Contraindications	A history of known hypersensitivity to Naloxone or any of its components		
Patient Education	Every person dispensed Naloxone under this standing order shall receive education regarding the risk factors of overdose, signs of an overdose, overdose response steps, and the use of Naloxone. Examples of educational materials that incorporate the above information may be found at http://www.naloxonesaves.org .		
Notification of Participation	Pharmacies choosing to participate in Naloxone dispensing under the authority of this standing order shall notify the Division of Public Health when initiating their participation; see directions for notification at http://www.naloxonesaves.org .		



 Elizabeth C. Tilson, MD, MPH
 National Provider ID: 1760540421

 March 12, 2018
 Date Signed

This order is effective immediately upon signing and may be revised or revoked by the State Health Director according to his/her discretion.